

174 x 1.5 oz (43g) Place Pacs CM1715

READY-TO-USE PLACE PACS  
**MEAL**



ACTIVE INGREDIENT: Warfarin Sodium Salt, 0.005%  
OTHER INGREDIENTS: 99.995%  
Contains Chlorinated Paraffin 100%  
KEEP OUT OF REACH OF CHILDREN  
**CAUTION**

**CONTRAC**  
RAT & MOUSE BAIT READY TO USE PLACE PACS

**KILLS RATS, MICE & MEADOW VOLE\***

*Kills Warfarin Resistant Norway Rats*

NET WT:  
**16.3 lbs (7.4kg)**

\*Kills Norway Rats, Norway Rats, Meadow Voles, House Mice, Brown Bore, and House Wren.

174 x 1.5 oz (43g) Place Pacs CM1715

**CONTRAC**  
RAT & MOUSE BAIT READY TO USE PLACE PACS



1 00 30835 40202 8

174 x 1.5 oz (43g) Place Pacs CM1715

**CONTRAC**  
RAT & MOUSE BAIT READY TO USE PLACE PACS

**KILLS RATS, MICE & MEADOW VOLE\***

*Kills Warfarin Resistant Norway Rats*

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STORAGE AND DISPOSAL**

**NET WEIGHT: 16.3 lbs (7.4kg)**

**174 x 1.5 oz (43g) Place Pacs**

**BELL LABORATORIES, INC.**

**MADE IN USA**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

**USA 1540 0102 0000 0000**

174 x 1.5 oz (43g) Place Pacs CM1715

**CONTRAC**  
RAT & MOUSE BAIT READY TO USE PLACE PACS



1 00 30835 40202 8

8 20204 58306 00 1



241CM-3



Corrugated Recycles

# 174 x 1.5 oz (43 g) Place Pacs

CM1715

READY-TO-USE PLACE PACS

# MEAL



ACTIVE INGREDIENT:  
Bromadiolone (CAS #28772-56-7): ..... 0.005%  
OTHER INGREDIENTS†: ..... 99.995%  
†Contains Denatonium Benzoate    TOTAL:    100.000%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See back panels for First Aid and additional precautionary statements.

# CONTRAC<sup>®</sup>

**RAT & MOUSE BAIT READY TO USE PLACE PACS**

## KILLS RATS, MICE & MEADOW VOLES\*

***Kills Warfarin Resistant Norway Rats***

NET WT:  
**16.3 lbs** (7.4 kg)

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

# 174 x 1.5 oz (43 g) Place Pacs

# CM1715

## FIRST AID

### HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

#### IF SWALLOWED:

- Call a poison control center, doctor, or 1-877-854-2494, or 1-800-858-7378\*\* immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

#### IF ON SKIN OR CLOTHING:

- Take off contaminated clothing. Rinse skin immediately with plenty of soap and water for 15–20 minutes. Call a poison control center or doctor for treatment advice.

#### IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15–20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

\*\* Also call this number for information on health concerns and pesticide incidents.

## NOTE TO PHYSICIAN

If swallowed or absorbed through the skin, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

## TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian at once.

## NOTE TO VETERINARIAN

**Anticoagulant Bromadiolone:** For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K<sub>1</sub>. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for sites, pests and application methods described on this label.

**IMPORTANT:** Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent accidents:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to huffed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

**Bait stations are mandatory for outdoor, above ground use. Tamper-resistant bait stations must be used wherever children, pets, non-target mammals, or birds may have access to the bait placement location.**

**USE RESTRICTIONS:** This product may only be used to control the following rodent pests in and around man-made structures: House mouse (*Mus musculus*), Norway rat (*Rattus norvegicus*), Roof rat (*Rattus rattus*), Cotton rat\* (*Sigmodon hispidus*), Eastern



## KILLS RATS, MICE & MEADOW VOLES\*

### Kills Warfarin Resistant Norway Rats

Norway rats, roof rats and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

ACTIVE INGREDIENT: Bromadiolone (CAS #28772-56-7):	0.005%
OTHER INGREDIENTS:	99.995%
Contains Denatonium Benzoate	TOTAL: 100.000%

## KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for First Aid and additional precautionary statements.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal:** Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

**Pesticide Container:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration. Call your Local Waste Agency for any questions on proper disposal.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



NET WEIGHT: 16.3 lbs (7.4 kg)  
174 x 1.5 oz (43 g) Place Pacs

EPA REG. NO. 12455-75

EPA EST. NO. 12455-WI-1

harvest mouse\* (*Reithrodontomys humuli*), Golden mouse\* (*Ochrotomys nuttalli*), Polynesian rat\* (*Rattus exulans*), Meadow vole\* (*Microtus pennsylvanicus*), White-throated woodrat\* (*Neotoma albigula*), Southern plains woodrat\* (*Neotoma micropus*), and Mexican woodrat\* (*Neotoma mexicana*). This product must be used in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

Burrow baiting with CONTRAC Rat & Mouse Bait Ready To Use Place Pacs is prohibited.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait. Do not use this product in sewers.

Do not sell this product in individual containers holding less than 16 pounds of bait.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats, mice or meadow voles\* will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

## APPLICATION DIRECTIONS:

**RATS:** Apply 2 to 10 Place Pacs at intervals of 15 to 30 feet per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease to appear.

**MICE AND MEADOW VOLES\*:** Place 1 Place Pac at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse or meadow vole\* activity cease to appear.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not get in eyes, on skin, or on clothing. Keep away from children, domestic animals and pets.

All handlers (including applicators) must wear long sleeved shirt and long pants, shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

## User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish, birds and other wildlife. Dogs, cats, and predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat


011116/08-15

**174 x 1.5 oz (43 g) Place Pacs**

**CM1715**

**CONTRAC<sup>®</sup>**

**RAT & MOUSE BAIT READY TO USE PLACE PACS**



1 00 30835 40202 8

 **Bell**  
LABORATORIES, INC.  
www.belllabs.com  
Madison, WI 53704 U.S.A.  
**MADE IN USA**

	<p><b>NOTE TO PHYSICIAN</b> If swallowed or absorbed through the skin, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.</p>				
	<p><b>TREATMENT FOR PET POISONING</b> If animal eats bait, call veterinarian at once.</p> <p><b>NOTE TO VETERINARIAN</b> <b>Anticoagulant Bromadiolone:</b> For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K<sub>1</sub>. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.</p> <p><b>DIRECTIONS FOR USE</b> It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See outer package label for complete DIRECTIONS FOR USE including USE RESTRICTIONS and BAITING INSTRUCTIONS. It is illegal to sell this product unless it is accompanied by a complete set of its accepted labeling.</p> <p><b>SEE LABEL ON OUTER PACKAGING FOR ADDITIONAL PRECAUTIONARY INFORMATION</b></p> <p><b>INDIVIDUAL SALE IS PROHIBITED BY LAW</b></p>		<p><b>CONTRAC<sup>®</sup></b> <b>RAT &amp; MOUSE BAIT READY TO USE PLACE PACS</b></p> <p><b>KILLS RATS, MICE &amp; MEADOW VOLES*</b></p> <p><b>Kills Warfarin Resistant Norway Rats</b> Norway rats, roof rats and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.</p> <p>ACTIVE INGREDIENT: Bromadiolone (CAS #28772-56-7): ..... 0.005% OTHER INGREDIENTS†: ..... 99.995% †Contains Denatonium Benzoate TOTAL 100.000%</p> <p><b>KEEP OUT OF REACH OF CHILDREN CAUTION</b></p> <p><b>SEE LABEL ON OUTER PACKAGING FOR ADDITIONAL PRECAUTIONARY INFORMATION</b> See back panels for First Aid and additional precautionary statements.</p> <p><b>NET WEIGHT: 1.5 oz (43 g)</b></p> <p>EPA REG. NO.: 12455-75 EPA EST. NO.: 12455-WI-1</p> <p>*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat</p> <p>Manufactured by:  <b>Bell</b> LABORATORIES, INC.</p> <p>3699 Kinsman Blvd. Madison, WI 53704 U.S.A. www.belllabs.com <b>MADE IN USA</b></p>		<p><b>FIRST AID</b> <b>HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE IF SWALLOWED:</b></p> <ul style="list-style-type: none"> <li>Call a poison control center, doctor, or 1-877-854-2494, or 1-800-858-7378** immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul> <p><b>IF ON SKIN OR CLOTHING:</b></p> <ul style="list-style-type: none"> <li>Take off contaminated clothing. Rinse skin immediately with plenty of soap and water for 15–20 minutes. Call a poison control center or doctor for treatment advice.</li> </ul> <p><b>IF IN EYES:</b></p> <ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15–20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul> <p>** Also call this number for information on health concerns and pesticide incidents.</p>
	>2006-7				111315/08-15